arXiv:1512.07739v1 [math.GN] 24 Dec 2015

REMARKS ON ASYMPTOTIC POWER DIMENSION

JACEK KUCAB AND MICHAEL ZARICHNYI

Abstract. Using a result of Dranishnikov and Smith we prove that, under some conditions, the asymptotic power dimension of a proper metric space coincides with the dimension of its subpower corona.

1. Introduction

The asymptotic power dimension is defined in [2]. In this note we use one result of [2] in order to derive the coincidence of the asymptotic power dimension of a proper metric space and the dimension of the subpower corona (this corona is introduced in [3]) of this space.

2. Preliminaries

A metric space X is called proper if every closed ball in X is compact.

Let $D > 0$. For a metric space (X, ρ) and a family $\mathcal{U}$ of subsets of X we say that $\mathcal{U}$ is D-bounded if for any $U \in \mathcal{U}$ we have $\operatorname{diam}(U) \leq D$; $\mathcal{U}$ is D-disjoint if for any $U, V \in \mathcal{U}$, $U \neq V$ we have $\inf\{\rho(x, y) | x \in U, y \in V\} > D$;

Recall that a metric space (X, ρ) is called cocompact if there is a compact subset $K \subset X$ such that $X = \cup\{f(K) \mid f \in \operatorname{Isom}(X)\}$, where $\operatorname{Isom}(X)$ is the set of all isometries of X.

A metric space (X, ρ) is said to be M-connected, for some $M > 0$ if for every $x, y \in X$ there exist $x = x_0, x_1, \ldots, x_k = y$ such that $\{x_0, x_1, \ldots, x_k\} \subset X$ and $\rho(x_i, x_{i+1}) \leq M$ for every $i = 0, 1, \ldots, k-1$.

Definition 2.1. For a metric space (X, ρ) and a nonnegative integer n we say that the asymptotic Assouad-Nagata dimension of X does not exceed n ($\operatorname{asdim}_{AN} X \leq n$) if there are constants $c > 0$ and $r_0 > 0$ such that for any $r > r_0$ there exist r-disjoint and cr-bounded families $\mathcal{U}_0, \mathcal{U}_1, \ldots, \mathcal{U}_n$ of subsets of X such that $\bigcup_{i=0}^{n} \mathcal{U}_i$ is a cover of X.

In a similar manner, one can introduce the notion of the asymptotic power dimension of a metric space as follows [2]:

Definition 2.2. For a metric space (X, ρ) and a nonnegative integer n we say that the asymptotic power dimension of X does not exceed n ($\operatorname{asdim}_P X \leq n$) if there are constants $\alpha > 0$ and $r_0 > 0$ such that for any $r > r_0$ there exist r-disjoint and r^α-bounded families $\mathcal{U}_0, \mathcal{U}_1, ..., \mathcal{U}_n$ of subsets of X such that $\bigcup_{i=0}^{n} \mathcal{U}_i$ is a cover of X.

Date: December 25, 2015.

2010 *Mathematics Subject Classification.* 54F45, 54E35.

Key words and phrases. Asymptotic dimension, Assouad-Nagata dimension, asymptotic power dimension, sublinear corona.

As usual we will say that $\operatorname{asdim}_P X = n$, for $n \geq 1$, when $\operatorname{asdim}_P X \leq n$ and it is not true that $\operatorname{asdim}_P X \leq n-1$.

It is well-known that for any metric ρ on a set X the function $\rho' : X \times X \to \mathbb{R}$, $\rho'(x,y) = \ln(1+\rho(x,y))$, is also a metric on X. In the sequel, we will denote (X,ρ) by X and (X,ρ') by X'. Note that for a proper and unbounded metric space X, the space X' is also proper and unbounded.

The following result is proved in [2]; we provide the proof for the reader's convenience.

Theorem 2.3. *For any metric space (X,ρ) we have $\operatorname{asdim}_P X = \operatorname{asdim}_{AN} X'$*

Proof. Let $\operatorname{asdim}_P X \leq n$. Let α and r_0 be as in the definition. Let $c = 2\alpha$ and let $r_0' = \max(r_0, \frac{\ln 2}{\alpha})$. Let $r > r_0'$. Then $e^r - 1 \geq r > r_0' \geq r_0$ and thus there exist (e^r-1)-disjoint(ρ) and $(e^r-1)^\alpha$-bounded(ρ) families $\mathcal{U}_0, \mathcal{U}_1, \ldots, \mathcal{U}_n$ of subsets of X such that $\bigcup_{i=0}^n \mathcal{U}_i$ is a cover of X. Let $x \in U$, $y \in V$ for any $U, V \in \mathcal{U}_i$, $U \neq V$, and any $i \in \{0,1,\ldots,n\}$. Then

$$\rho'(x,y) = \ln(1+\rho(x,y)) > \ln(1+e^r-1) = r.$$

Let now $x, y \in U$ for any $U \in \mathcal{U}_i$ and any $i \in \{0,1,\ldots,n\}$. Then we have

$$\rho'(x,y) = \ln(1+\rho(x,y)) \leq \ln(1+(e^r-1)^\alpha) \leq \ln(1+e^{\alpha r}) \leq \ln(2e^{\alpha r}) < \ln e^{2\alpha r} = cr,$$

for $\alpha r > \ln 2$, i.e. $e^{\alpha r} > 2$. Therefore for any $i \in \{0,1,\ldots,n\}$, the family $\mathcal{U}_i$ is r-disjoint(ρ') and cr-bounded(ρ'). Hence $\operatorname{asdim}_{AN} X' \leq n$ and therefore $\operatorname{asdim}_{AN} X' \leq \operatorname{asdim}_P X$.

Let now $\operatorname{asdim}_{AN} X' \leq n$. Let c and r_0' be as in the definition. Let $\alpha = c + 2^c$ and $r_0 = e^{r_0'}$. Let $r > r_0$. Hence $r+1 > r_0 > 1$, i.e. $\ln(r+1) > r_0'$. Then there exist $\ln(r+1)$-disjoint(ρ') and $c\ln(r+1)$-bounded(ρ') families $\mathcal{U}_0, \mathcal{U}_1, \ldots, \mathcal{U}_n$ of subsets of X such that $\bigcup_{i=0}^n \mathcal{U}_i$ is a cover of X. Let $x \in U$, $y \in V$ for any $U, V \in \mathcal{U}_i$, $U \neq V$, and any $i \in \{0,1,\ldots,n\}$. Then

$$\rho(x,y) = e^{\rho'(x,y)} - 1 > e^{\ln(r+1)} - 1 = r.$$

Let now $x, y \in U$ for any $U \in \mathcal{U}_i$ and any $i \in \{0,1,\ldots,n\}$. Then

$$\rho(x,y) = e^{\rho'(x,y)} - 1 \leq e^{c\ln(r+1)} - 1 < (r+1)^c < (2r)^c < r^\alpha,$$

since for any r we have $\alpha > c + \log_r 2^c$ and hence $r^\alpha > r^c 2^c$. Therefore for any $i \in \{0,1,\ldots,n\}$ the family $\mathcal{U}_i$ is r-disjoint(ρ) and r^α-bounded(ρ). Therefore $\operatorname{asdim}_P X \leq n$. We conclude that $\operatorname{asdim}_P X \leq \operatorname{asdim}_{AN} X'$ and therefore $\operatorname{asdim}_P X = \operatorname{asdim}_{AN} X'$. □

The notions of sublinear (resp. subpower) corona are introduced in [1] and [3] respectively. The necessary definitions are given below.

Definition 2.4. We recall that a function $p : \mathbb{R}_+ \to \mathbb{R}_+$ is called asymtotically subpower (resp. asymtotically sublinear) if for any $\alpha > 0$ [$c > 0$] there exists $r_0 > 0$ such that for any $x > r_0$ statement $p(x) < x^\alpha$ (resp. $p(x) < cx$) holds.

Definition 2.5. Let (X,ρ) be an unbounded proper metric space with a basepoint x_0. We say that a continous, bounded function $f : X \to \mathbb{R}$ is Higson subpower (resp. Higson sublinear) if for any asymtotically subpower function p (resp. asymtotically sublinear function p) we have $\lim_{|x|\to\infty} \operatorname{diam}(f(B_{p(|x|)}(x))) = 0$, where $|x| = \rho(x, x_0)$.

It is not hard to see that Higson subpower functions as well as Higson sublinear functions on a proper, unbounded metric space form subalgebras of algebra of all continuous and bounded functions on X. We will denote them by $CB_P(X)$ and $CB_L(X)$ respectively.

Theorem 2.6. *For an unbounded, proper metric space (X, ρ) we have $CB_P(X) = CB_L(X')$.*

Proof. Let (X, ρ) be an unbounded proper metric space with a basepoint x_0.

First we are going to prove that $CB_P(X) \subset CB_L(X')$. Convention: we use $B'_r(x)$ for the balls in X' and also $|x|' = \rho(x, x_0)$.

Let $f \in CB_P(X)$ and let $\phi\colon \mathbb{R}_+ \to \mathbb{R}_+$ be any asymptotically sublinear function. Note that

$$\begin{aligned} B'_{\phi(|x|')}(x) =& \{y \in X | \rho'(y, x) < \phi(|x|')\} \\ =& \{y \in X \mid \ln(1 + \rho(y, x)) < \phi(\ln(1 + |x|))\} \\ =& \{y \in X \mid \rho(y, x) < e^{\phi(\ln(1+|x|))} - 1\}. \end{aligned}$$

Denoting by $\psi(|x|) = e^{\phi(\ln(1+|x|))} - 1$, the last set is just $B_{\psi(|x|)}(x)$. Now for every $n \in \mathbb{N}$ there is some $c_n \in \mathbb{R}$ that

$$\ln \psi(|x|) < \ln e^{\phi(\ln(1+|x|))} = \phi(\ln(1 + |x|)) < \frac{1}{n} \ln(1 + |x|)),$$

whenever $\ln(1 + |x|)) > c_n$ (or equivalently $|x| > e^{c_n} - 1$). The latter means that for every $n \in \mathbb{N}$ there is $c_n \in \mathbb{R}$ such that $\psi(|x|) < (1 + |x|)^{1/n} < |x|^{2/n}$ whenever $|x| > \max(2, e^{c_n} - 1)$ and so ψ is an asymptotically subpower function (assuming by definition $\psi(\lambda) = 1$ for those $\lambda \in \mathbb{R}_+$ for which there is no $x \in X$, such that $\lambda = |x|$). Thus $\operatorname{diam}(f(B'_{\phi(|x|')}(x))) = \operatorname{diam}(f(B_{\psi(|x|)}(x))) \to 0$. We conclude that $f \in CB_L(X')$.

Now we are going to prove that $CB_L(X') \subset CB_P(X)$.

Let $f \in CB_L(X')$ and let $\phi\colon \mathbb{R}_+ \to \mathbb{R}_+$ be an asymptotically subpower function. Note that $\phi + 1$ is an asymptotically subpower function as well. Let $c_1 > 0$ be large enough, such that $1 + \phi(x) < x < x + 1$ for every $x > c_1$ and for $n \in \mathbb{N}$, $n > 1$, let $c_n > c_{n-1}$ be large enough, such that $1 + \phi(x) < x^{1/n} < (1 + x)^{1/n}$ for every $x > c_n$. We will now construct a function $\psi\colon \mathbb{R}_+ \to \mathbb{R}_+$ as follows: $\psi(x) = x/n$ for $x \in (c_n, c_{n+1}]$, $n = 1, 2, \ldots$, and $\psi(x) = 1$ for $x \in (0, c_1]$. The fact that ψ is an asymptotically sublinear function is obvious, as for every $n \in \mathbb{N}$ there exists c_{n+1} such that for every $x > c_{n+1}$ we have $\psi(x) \leq \frac{1}{n+1} x < \frac{1}{n} x$. We will now show that for $x \in X$ with $|x|$ large enough, $B_{\phi(|x|)}(x) \subset B'_{\psi(|x|')}(x)$. Let then $y \in B_{\phi(|x|)}(x)$. Therefore $\rho(x, y) < \phi(|x|)$. Hence

$$\rho'(x, y) = \ln(1 + \rho(x, y)) < \ln(1 + \phi(|x|)) < \ln(1 + |x|)^{1/n} = \frac{1}{n} \ln(1 + |x|)$$

whenever $|x| > c_n$, hence $\rho'(x, y) < \psi(|x|')$ and $y \in B'_{\psi(|x|')}(x)$ for $|x|$ large enough.

Since $\operatorname{diam}(f(B'_{\psi(|x|')}(x)) \to 0$, we see that $\operatorname{diam}(f(B_{\phi(|x|)}(x))$ and hence $f \in CB_P(X)$. □

Let $h_P X$ and $h_L X$ denote the compactifications of X with respect to the algebras $CB_P(X)$ and $CB_L(X)$ respectively. Let $\nu_P X$ and $\nu_L X$ denote the coronas of these compactifications.

Corollary 2.7. *For an unbounded proper metric space (X, ρ) we have $\dim \nu_P X = \dim \nu_L X'$.*

Proof. Let (X, ρ) be an unbounded proper metric space. By Theorem 2.6 we have $CB_P(X) = CB_L(X')$ and hence there is a homeomorphism $h_P X \to h_L X'$ which extends the identity and this homeomorphism induces a homeomorphism on the coronas. □

Lemma 2.8. *For a metric space (X, ρ) a set Γ of isometries of X coincides with the set Γ' of isometries of X'.*

Proof. For any $\gamma \in \Gamma$ and any $x, y \in X$ we have

$$\rho'(\gamma(x), \gamma(y)) = \ln(1 + \rho(\gamma(x), \gamma(y))) = \ln(1 + \rho(x, y)) = \rho'(x, y),$$

so $\Gamma \subset \Gamma'$ and for any $\gamma' \in \Gamma'$ and any $x, y \in X$ we have

$$\rho(\gamma'(x), \gamma'(y)) = e^{\rho'(\gamma'(x), \gamma'(y))} - 1 = e^{\rho'(x,y)} - 1 = \rho(x, y),$$

therefore $\Gamma' \subset \Gamma$, i.e. $\Gamma = \Gamma'$. □

Corollary 2.9. *If X is a cocompact metric space, so is X'.*

The following is proved in [1] (see Corollary 3.11 therein).

Theorem 2.10. *For a cocompact, proper, unbounded and M-connected (for some $M > 0$) metric space X with finite $\operatorname{asdim}_{AN} X$ we have*

$$\operatorname{asdim}_{AN} X = \dim \nu_L X$$

Corollary 2.11. *For a proper, unbounded, cocompact and M-connected (for some $M > 0$) metric space X with finite $\operatorname{asdim}_P X$ we have*

$$\operatorname{asdim}_P X = \dim \nu_P X.$$

Proof. Let (X, ρ) be a proper, unbounded, cocompact and M-connected (for some $M > 0$) metric space with finite $\operatorname{asdim}_P X$. Then X' is proper, unbounded and cocompact with finite $\operatorname{asdim}_{AN} X'$. Since for any $x, y \in X$ we have $\rho'(x, y) = \ln(1 + \rho(x, y)) \leq \rho(x, y)$, the space X' is M-connected as well. Using previous results we have:

$$\operatorname{asdim}_P X = \operatorname{asdim}_{AN} X' = \dim \nu_L X' = \dim \nu_P X$$

□

Corollary 2.12. *For a finitely generated group G with word metric and of finite subpower asymptotic dimension we have:*

$$\operatorname{asdim}_P G = \dim \nu_P G.$$

3. Remarks

Corollary 2.11 provides an answer to one question formulated in [2].

The remetrization $\rho \mapsto \rho'$ can be used also for reducing some questions concerning the asymptotic power dimension to the corresponding questions concerning the asymptotic Assouad-Nagata dimension. In particular, we conjecture that this remetrization would be useful in finding of an alternative proof of a characterization of the asymptotic power dimension in terms of mappings into polyhedra (this was obtained by the first author).

References

[1] A.N. Dranishnikov, J. Smith, On asymptotic Assouad-Nagata dimension, Topology and its Applications, Volume 154, Issue 4, 2007, 934–952.
[2] J. Kucab, M. Zarichnyi, On asymptotic power dimension, Topology and its Applications (in press).
[3] J. Kucab, M. Zarichnyi, Subpower Higson corona of a metric space, Algebra Discrete Math., 17:2 (2014), 280–287.

Faculty of Mathematics and Natural Sciences, University of Rzeszów, Rejtana 16 A, 35-310 Rzeszów, Poland
E-mail address: mzar@litech.lviv.ua

Department of Mechanics and Mathematics, Lviv National University, Universytetska Str. 1, 79000 Lviv, Ukraine

Faculty of Mathematics and Natural Sciences, University of Rzeszów, Rejtana 16 A, 35-310 Rzeszów, Poland
E-mail address: mzar@litech.lviv.ua